

# KAZHDAN GROUPS WHOSE FC-RADICAL IS NOT VIRTUALLY ABELIAN

MIKHAIL ERSHOV

## 1. INTRODUCTION

If  $G$  is a group, the *FC-radical* of  $G$ , denoted by  $FC(G)$ , is the set of all elements of  $G$  which centralize a finite index subgroup of  $G$ . Equivalently,  $FC(G)$  is the set of all elements of  $G$  with finite conjugacy class.

In [PV] Popa and Vaes asked the following question:

**Question 1.1.** *Does there exist a residually finite (discrete) group with Kazhdan's property (T) whose FC-radical is not virtually abelian?*

This question was motivated by [PV, Theorem 6.4(b)], which asserts that any group satisfying these conditions admits a free ergodic profinite action whose associated  $\text{II}_1$  factor has all positive real numbers in its fundamental group. No Kazhdan group with the latter property was previously known.

In this short note we give a positive answer to Question 1.1 using Golod-Shafarevich groups.

We shall prove the following theorem:

**Theorem 1.2.** *Every Golod-Shafarevich group has a residually finite quotient whose FC-radical is not virtually abelian.*

In [Er] it was shown that there exist Golod-Shafarevich groups with property (T). Since property (T) is preserved by quotients, applying Theorem 1.2 to any Golod-Shafarevich group with (T), we obtain a group which settles Question 1.1.

**Acknowledgements.** I am grateful to Mark Sapir for telling me about Question 1.1.

## 2. CONSTRUCTION

Informally speaking, a finitely generated group  $G$  is Golod-Shafarevich if it has a presentation with a “small” set of relators, where relators are counted with suitable weights. The formal definition is given below.

**Definition.** Let  $G$  be a finitely generated group. Given a prime  $p$  let  $\mathbb{F}_p[G]$  be the  $\mathbb{F}_p$ -group algebra of  $G$  and  $I$  the augmentation ideal of  $\mathbb{F}_p[G]$ . Let  $\{\omega_n G\}_{n \geq 1}$  be the Zassenhaus  $p$ -filtration of  $G$  defined by  $\omega_n G = \{g \in G : g - 1 \in I^n\}$ . For each  $g \in G \setminus \bigcap_{n \in \mathbb{N}} \omega_n G$  we put  $\deg_p(g)$  to be the largest  $n$  such that  $g \in \omega_n G$ .

It is well known that the subgroups  $\{\omega_n G\}$  are of finite index in  $G$ . Moreover, they form a base for the pro- $p$  topology on  $G$ , and thus  $\bigcap_{n \in \mathbb{N}} \omega_n G$  is the kernel of the natural map from  $G$  to its pro- $p$  completion  $G_{\hat{p}}$ . In particular, if  $G$  is residually- $p$ , then  $\bigcap_{n \in \mathbb{N}} \omega_n G = \{1\}$ , so  $\deg_p(g)$  is defined for any  $g \in G \setminus \{1\}$ .

---

The author is supported by the NSF grant DMS-0901703.

**Definition.** Fix a prime number  $p$ .

- (a) A group presentation  $\langle X|R \rangle$ , with  $X$  finite, is said to satisfy the *Golod-Shafarevich* (GS) condition (with respect to  $p$ ), if there is a real number  $t \in (0, 1)$  such that

$$1 - H_X(t) + H_R(t) < 0 \text{ where } H_X(t) = |X|t \text{ and } H_R(t) = \sum_{r \in R} t^{\deg_p(r)}.$$

- (b) A group  $G$  is called *Golod-Shafarevich* if it has a presentation satisfying the Golod-Shafarevich condition.

Any Golod-Shafarevich group  $G$  is infinite. In fact, its pro- $p$  completion  $G_{\hat{p}}$  must be infinite and moreover satisfies a number of largeness properties (see, e.g., [Ze], [Er] and references therein for precise statements). We shall only use a very weak statement about Golod-Shafarevich groups:

**Proposition 2.1.** *If a group  $G$  is Golod-Shafarevich with respect to  $p$ , then its pro- $p$  completion  $G_{\hat{p}}$  is not virtually abelian.*

*Proof of Theorem 1.2.* Let  $G$  be a Golod-Shafarevich group, so that  $G$  has a presentation  $\langle X|R \rangle$  with  $1 - H_X(t) + H_R(t) < 0$  for some  $t \in (0, 1)$ , and let  $\varepsilon = -(1 - H_X(t) + H_R(t))$ .

Let  $k_0 \in \mathbb{N}$  be such that  $t^{k_0} < \frac{\varepsilon}{8}$ . By Proposition 2.1 we can choose  $x_1, y_1 \in \omega_{k_0}G$  which do not commute in  $G_{\hat{p}}$ . Then there exists  $k_1 > k_0$  such that  $x_1$  and  $y_1$  do not commute in  $G/\omega_{k_1}G$ . By making  $k_1$  larger we can also assume that  $t^{k_1} < \frac{\varepsilon}{16}$ .

Let  $S_1$  be a finite generating set for  $\omega_{k_1}G$ , let  $R_1 = \{[x_1, s], [y_1, s] : s \in S_1\}$  and  $G_1 = G/\langle R_1 \rangle^G$ . Note that if  $\bar{x}_1$  and  $\bar{y}_1$  are the images of  $x_1$  and  $y_1$  in  $G_1$ , then

- (i)  $G/\omega_{k_1}G \cong G_1/\omega_{k_1}G_1$ , so  $\bar{x}_1$  and  $\bar{y}_1$  do not commute modulo  $\omega_{k_1}G_1$ ;
- (ii)  $\bar{x}_1$  and  $\bar{y}_1$  lie in the FC-radical of  $G_1$  (and the same is true for any quotient of  $G_1$ ).

The group  $G_1$  need not be Golod-Shafarevich, but it surjects onto the group  $\widehat{G}_1 = G_1/\langle x_1, y_1 \rangle^{G_1}$  which is Golod-Shafarevich by construction.

Thus, the pro- $p$  completion of  $G_1$  is not virtually abelian, so we can find elements  $x_2, y_2 \in \omega_{k_1}G$  and  $k_2 > k_1$  such that  $x_2$  and  $y_2$  do not commute in  $G_2/\omega_{k_2}G_2$  and  $t^{k_2} < \frac{\varepsilon}{32}$ .

Let  $S_2$  be a finite generating set for  $\omega_{k_2}G$ , let  $R_2 = \{[x_2, s], [y_2, s] : s \in S_2\}$  and  $G_2 = G/\langle R_1 \cup R_2 \rangle^G$ . By construction we have  $G_1/\omega_{k_2}G_1 \cong G_2/\omega_{k_2}G_2$ , the images of  $x_2$  and  $y_2$  in  $G_2$  lie in the FC-radical of  $G_2$ , and  $G_2$  surjects onto a Golod-Shafarevich group.

Continuing this process indefinitely we obtain a sequence of groups  $G = G_0 \rightarrow G_1 \rightarrow G_2 \rightarrow \dots$ , elements  $\{x_i, y_i\}_{i \in \mathbb{N}}$  of  $G$  and integers  $k_0 < k_1 < k_2 < \dots$  s.t.

- (i)  $G_{i+1}$  is a quotient of  $G_i$  for all  $i$
- (ii)  $G_i$  surjects onto the group  $G/\langle \cup_{j=1}^i \{x_j, y_j\} \rangle^G$
- (iii)  $x_i$  and  $y_i$  lie in  $\omega_{k_{i-1}}G$ , and  $t^{k_{i-1}} < \frac{\varepsilon}{2^{i+2}}$
- (iv) The images of  $x_i$  and  $y_i$  in  $G_{i-1}/\omega_{k_i}G_{i-1}$  do not commute
- (v)  $G_{i-1}/\omega_{k_i}G_{i-1} \cong G_j/\omega_{k_i}G_j$  for all  $j \geq i$
- (vi) The images of  $x_i$  and  $y_i$  in  $G_i$  lie in the FC-radical of  $G_i$

Now let  $G_{\infty}$  be the inductive limit of  $\{G_i\}$ ; in other words, if  $G_i = G/N_i$ , we let  $N_{\infty} = \cup_{i \in \mathbb{N}} N_i$  and  $G_{\infty} = G/N_{\infty}$ . Let  $Q$  be the image of  $G_{\infty}$  in its pro- $p$  completion, that is,  $Q = G_{\infty}/\cap_{n \in \mathbb{N}} \omega_n G_{\infty}$ .

Condition (ii) implies that  $G$  surjects onto the group  $G/\langle \cup_{j=1}^{\infty} \{x_j, y_j\} \rangle^G$  which is Golod-Shafarevich by (iii). Thus, by Proposition 2.1 the group  $Q$  is infinite. Since  $Q$  is a subset of  $(G_{\infty})_{\hat{p}}$ , it is also residually finite.

Let  $\pi : G \rightarrow Q$  be the natural projection. By condition (vi) the FC-radical of  $Q$  contains the subgroup  $H$  generated by the elements  $\{\pi(x_i), \pi(y_i)\}_{i \in \mathbb{N}}$ . It remains to show that  $H$  is not virtually abelian. Suppose not, so  $H$  contains a finite index abelian subgroup  $A$ . Then there exists integers  $i < j$  such that  $\pi(x_i x_j^{-1}) \in A$  and  $\pi(y_i y_j^{-1}) \in A$ . Conditions (iv) and (v) imply that  $\pi(x_i)$  and  $\pi(y_i)$  do not commute modulo  $\omega_{k_i} Q$ . Thus, if  $\varphi_i$  is the projection map  $Q \rightarrow Q/\omega_{k_i} Q$ , then  $\varphi_i \pi([x_i, y_i]) \neq 1$ . On the other hand, by construction  $x_j, y_j \in \omega_{k_i} G$ , so  $\varphi_i \pi(x_j) = \varphi_i \pi(y_j) = 1$ . Therefore,

$$\varphi_i \pi([x_i, y_i]) = \varphi_i \pi([x_i x_j^{-1}, y_i y_j^{-1}]) \in \varphi_i([A, A]) = \{1\}.$$

The obtained contradiction shows that the FC-radical of  $Q$  is not virtually abelian, which finishes the proof.  $\square$

#### REFERENCES

- [Er] M. Ershov, *Golod-Shafarevich groups with property (T) and Kac-Moody groups*, Duke Math. J. 145 (2008), no. 2, 309–339
- [PV] S. Popa and S. Vaes, *On the fundamental group of  $\text{II}_1$  factors and equivalence relations arising from group actions*, arXiv:0810.0706, to appear in the Proceedings of the Conference in Honor of Alain Connes' 60th birthday, "Non-Commutative Geometry" April 2-6, 2007, IHP Paris.
- [Ze] E. Zelmanov, *Infinite algebras and pro-p groups*. Infinite groups: geometric, combinatorial and dynamical aspects, 403–413, Progr. Math., 248, Birkhäuser, Basel, 2005.

UNIVERSITY OF VIRGINIA

*E-mail address:* `ershov@virginia.edu`